

Coalition and Group Announcement Logic

Rustam Galimullin

Natasha Alechina

School of Computer Science
University of Nottingham
Nottingham, UK

{rustam.galimullin, natasha.alechina}@nottingham.ac.uk

Dynamic epistemic logics which model abilities of agents to make various announcements and influence each other's knowledge have been studied extensively in recent years. Two notable examples of such logics are Group Announcement Logic and Coalition Announcement Logic. They allow us to reason about what groups of agents can achieve through joint announcements in non-competitive and competitive environments. In this paper, we consider a combination of these logics – Coalition and Group Announcement Logic and provide its complete axiomatisation. Moreover, we partially answer the question of how group and coalition announcement operators interact, and settle some other open problems.

1 Introduction

To introduce the logics we will be working with in this paper, we start with an example loosely based on the one from [16]. Let us imagine that Ann, Bob, and Cath are travelling by train from Nottingham to Liverpool through Manchester. Cath was sound asleep all the way, and she has just woken up. She does not know whether the train passed Manchester, but Ann and Bob know that it has not. Now, if the train driver announces that the train is approaching Manchester, then Cath, as well as Ann and Bob, knows that they have not passed the city yet. To reason about changes in agents' knowledge after public announcements, we can use Public Announcement Logic (**PAL**) [15]. Returning to the example, let us assume that the train driver does not announce anything, so that Cath is not aware of her whereabouts. Ann and Bob may tell her whether they passed Manchester. In other words, Ann and Bob have an announcement that can influence Cath's knowledge. An extension of **PAL**, Group Announcement Logic (**GAL**) [2], deals with the *existence* of announcements by groups of agents that can achieve certain results. Now, let us assume that Ann does not want to disclose to Cath their whereabouts and Bob does, i.e. Ann and Bob have different goals. Then, it is clear that no matter what Ann says, the coalition of Bob and Cath can achieve the goal of Cath knowing that the train has not passed Manchester, that is, Bob can communicate this information to Cath. On the other hand, if Ann and Bob work together, then they have an announcement (for example, a tautology 'It either rains in Liverpool or it doesn't'), such that whatever Cath says, she remains unaware of her whereabouts. For this type of strategic behaviour, another extension of **PAL** – Coalition Announcement Logic (**CAL**) – has been introduced in [3].

CAL joins two logical traditions: Dynamic Epistemic Logic, of which **PAL** is a representative, and Coalition Logic (**CL**) [14]. The latter allows us to reason about whether a coalition of agents has a strategy to achieve some goal, no matter what the agents outside of the coalition do. **CL** essentially talks about concurrent games, and the actions that the agents execute are arbitrary actions (strategies in one-shot games). So, from this perspective, **CAL** is a coalition logic with available actions restricted to public announcements.

To the best of our knowledge, there is no complete axiomatisation of **CAL** [3, 11, 4, 5] or any other logic with coalition announcement operators. In this paper, we consider Coalition and Group

Announcement Logic (**CoGAL**), a combination of **GAL** and **CAL**, which includes operators for both group and coalition announcements. The main result of this paper is a sound and complete axiomatisation of **CoGAL**. As part of this result, we study the interplay between group and coalition announcement operators, and partially settle the question on their interaction that was stated as an open problem in [11, 5].

2 Coalition and Group Announcement Logic

2.1 Syntax and Semantics

Throughout the paper, let a finite set of agents A , and a countable set of propositional variables P be given. The language of the logic is comprised of the language of classical propositional logic with added operators for agents' knowledge $K_a\varphi$ (reads 'agent a knows φ '), and public announcement $[\psi]\varphi$ (reads 'after public announcement that ψ , φ holds), group $[G]\varphi$ ('after any public announcement by group of agents G , φ holds), and coalition announcements $\llbracket G \rrbracket\varphi$ ('for every public announcement by coalition of agents G there is an announcement by other agents $A \setminus G$, such that φ holds after joint simultaneous announcement').

Definition 2.1. (Language) The *language of coalition and group announcement logic* \mathcal{L}_{CoGAL} is as follows:

$$\varphi, \psi ::= p \mid \neg\varphi \mid (\varphi \wedge \psi) \mid K_a\varphi \mid [\varphi]\psi \mid [G]\varphi \mid \llbracket G \rrbracket\varphi,$$

where $p \in P$, $a \in A$, $G \subseteq A$, and all the usual abbreviations of propositional logic (such as $\vee, \rightarrow, \leftrightarrow$) and conventions for deleting parentheses hold. The dual operators are defined as follows: $\widehat{K}_a\varphi \leftrightarrow \neg K_a\neg\varphi$, $\langle \varphi \rangle\psi \leftrightarrow \neg[\varphi]\neg\psi$, $\langle G \rangle\varphi \leftrightarrow \neg[G]\neg\varphi$, and $\llbracket G \rrbracket\varphi \leftrightarrow \neg\llbracket G \rrbracket\neg\varphi$. Observe that $\langle G \rangle\varphi$ means that G has an announcement after which φ holds, and $\llbracket G \rrbracket\varphi$ means that G has an announcement such that after it is made simultaneously with any announcement by $A \setminus G$, φ holds. The latter corresponds to the Coalition Logic operator, but for announcements instead of arbitrary actions.

We define \mathcal{L}_{GAL} as the language without the operator $\llbracket G \rrbracket$, \mathcal{L}_{PAL} the language without $[G]$ as well, and \mathcal{L}_{EL} the purely epistemic language which in addition does not contain announcement operators $[\varphi]$.

Next definition is needed for technical reasons in the formulation of infinite rules of inference in Definition 2.5. We want the rules to work for a class of different types of premises. Ultimately, we require premises to be expressions of depth n of the type $\varphi_1 \rightarrow \Box_1(\varphi_2 \rightarrow \dots (\varphi_n \rightarrow \Box_n\#) \dots)$, where \Box_i is either K_a or $[\psi]$ for some $a \in A$ and $\psi \in \mathcal{L}_{CoGAL}$, atom $\#$ denotes a placement of a formula to which a derivation is applied, and some φ 's and \Box 's can be omitted. This condition is captured succinctly by necessity forms originally introduced by Goldblatt in [13].

Definition 2.2. (Necessity forms) Let $\varphi \in \mathcal{L}_{CoGAL}$, then *necessity forms* [13] are inductively defined as follows:

$$\eta ::= \# \mid \varphi \rightarrow \eta(\#) \mid K_a\eta(\#) \mid [\varphi]\eta(\#).$$

The atom $\#$ has a unique occurrence in each necessity form. The result of the replacement of $\#$ with φ in some $\eta(\#)$ is denoted as $\eta(\varphi)$.

Whereas formulas of coalition logic [14] are interpreted in game structures, formulas of **CoGAL** are interpreted in epistemic models. Let us consider an example of such a model first. In Figure 1 there are three agents: a (Ann), b (Bob), and c (Cath). Let p denote the proposition that 'The train has passed Manchester.' There are two states in the model M : a state w where $\neg p$ is true, and a state v where p is




Figure 1: Train example

true; and only one state in model M^{-p} which denotes M updated by the announcement $\neg p$ (the process of updating the model is described below). Let the w be the actual state. Edges connect states that an agent cannot distinguish. In the actual state w of M , Cath (agent c) does not know whether p is true. Ann and Bob, on the contrary, know that p is false. Now suppose that Bob announces that $\neg p$. This truthful public announcement ‘deletes’ all the states where p is true, and the corresponding epistemic indistinguishability relations; in this example, v is ‘deleted,’ and the resulting model is M^{-p} . After this announcement Cath knows $\neg p$, or, formally, $[\neg p]K_c\neg p$. In this paper, within group and coalition announcements, we only quantify over announcements of formulas of the type $K_a\phi$. If a group consists only of Cath, who does not know $\neg p$ and hence cannot announce $K_c\neg p$, the following holds in state w of M : $[c](\neg K_c\neg p \wedge \neg K_cp)$, i.e. whatever c announces, she still does not know whether p after the announcement¹. Also, Ann and Bob can remain silent (or announce a tautology \top) and preclude Cath from knowing that $\neg p$. In other words, there is announcement by their group such that after it is made, agent c does not know the value of p : $\langle\{a, b\}\rangle(\neg K_c\neg p \wedge \neg K_cp)$. Moreover, this holds whatever Cath announces at the same time: $\langle\{\{a, b\}\}\rangle(\neg K_c\neg p \wedge \neg K_cp)$. On the other hand, a coalition consisting of Ann and Cath does not have such a power, since Bob can always announce that $\neg p$: $\neg\langle\{a, c\}\rangle(\neg K_c\neg p \wedge \neg K_cp)$, or, equally, $[\langle\{a, c\}\rangle](K_c\neg p \vee K_cp)$.

Now, we provide formal definitions.

Definition 2.3. (Epistemic model) An *epistemic model* is a triple $M = (W, \sim, V)$, where

- W is a non-empty set of states;
- $\sim: A \rightarrow \mathcal{P}(W \times W)$ assigns an equivalence relation to each agent; we will denote relation assigned to agent $a \in A$ by \sim_a ;
- $V: P \rightarrow \mathcal{P}(W)$ assigns a set of states to each propositional variable.

A pair (W, \sim) is called an *epistemic frame*, and a pair (M, w) with $w \in W$ is called a *pointed model*. An announcement in a pointed model (M, w) results in an *updated pointed model* (M^ϕ, w) . Here $M^\phi = (W^\phi, \sim^\phi, V^\phi)$, and $W^\phi = \llbracket \phi \rrbracket_M$, $\sim_a^\phi = \sim_a \cap (\llbracket \phi \rrbracket_M \times \llbracket \phi \rrbracket_M)$, and $V^\phi(p) = V(p) \cap \llbracket \phi \rrbracket_M$. Generally speaking, an updated pointed model (M^ϕ, w) is a restriction of the original one to the states where ϕ holds.

Let \mathcal{L}_{EL}^G denote the set of formulas of the type $\bigwedge_{i \in G} K_i \phi_i$, where for every $i \in G$ it holds that $\phi_i \in \mathcal{L}_{EL}$. These are the formulas we will be quantifying over in modalities of the form $[G]$ and $\langle\langle G \rangle\rangle$.

Definition 2.4. (Semantics) Let a pointed model (M, w) with $M = (W, \sim, V)$, $a \in A$, and $\phi, \psi \in \mathcal{L}_{CoGAL}$ be given.

$$\begin{aligned}
(M, w) \models p & \quad \text{iff} \quad w \in V(p) \\
(M, w) \models \neg\phi & \quad \text{iff} \quad (M, w) \not\models \phi \\
(M, w) \models \phi \wedge \psi & \quad \text{iff} \quad (M, w) \models \phi \text{ and } (M, w) \models \psi \\
(M, w) \models K_a\phi & \quad \text{iff} \quad \forall v \in W : w \sim_a v \text{ implies } (M, v) \models \phi \\
(M, w) \models [\phi]\psi & \quad \text{iff} \quad (M, w) \models \phi \text{ implies } (M^\phi, w) \models \psi \\
(M, w) \models [G]\phi & \quad \text{iff} \quad \forall \psi \in \mathcal{L}_{EL}^G : (M, w) \models [\psi]\phi \\
(M, w) \models \langle\langle G \rangle\rangle\phi & \quad \text{iff} \quad \forall \psi \in \mathcal{L}_{EL}^G \exists \chi \in \mathcal{L}_{EL}^{A \setminus G} : (M, w) \models \psi \rightarrow \langle\psi \wedge \chi\rangle\phi
\end{aligned}$$

¹For readability, we use $[c]$ rather than $[\{c\}]$ for singleton coalitions.

Formula φ is called *valid* if for any pointed model (M, w) it holds that $(M, w) \models \varphi$.

The semantics for the ‘diamond’ versions of knowledge, public and group announcement operators ($\widehat{K}_a\varphi$, $\langle\varphi\rangle\psi$, and $\langle G\rangle\varphi$ respectively) are obtained by changing \forall to \exists and ‘implies’ to ‘and’ in the corresponding lines. The semantics for a dual of the coalition announcement operator is as follows:

$$(M, w) \models \langle[G]\rangle\varphi \quad \text{iff} \quad \exists\psi \in \mathcal{L}_{EL}^G \forall\chi \in \mathcal{L}_{EL}^{A \setminus G} : (M, w) \models \psi \wedge [\psi \wedge \chi]\varphi,$$

which corresponds to ‘there is an announcement by agents from G , such that whatever other agents $A \setminus G$ announce at the same time, φ holds.’

Note that following [8, 7, 2, 3, 5, 9, 11, 4] we restrict formulas which agents in a group or coalition can announce to formulas of \mathcal{L}_{EL} . This allows us to avoid circularity in the definition.

2.2 Axiomatisation and Some Logical Properties

In this section we present an axiomatisation of **CoGAL** and show its soundness.

Definition 2.5. *Axiomatisation* of **CoGAL** is a union of axiomatisation of **GAL** [2], interaction axiom for group and coalition announcements A11, rule of inference for coalition announcements R6, and necessitation R4.

- | | |
|---|--|
| (A0) instantiations of propositional tautologies, | (A10) $[G]\varphi \rightarrow [\psi]\varphi$, where $\psi \in \mathcal{L}_{EL}^G$, |
| (A1) $K_a(\varphi \rightarrow \psi) \rightarrow (K_a\varphi \rightarrow K_a\psi)$, | (A11) $\langle[G]\rangle\varphi \rightarrow \langle G\rangle[A \setminus G]\varphi$, |
| (A2) $K_a\varphi \rightarrow \varphi$, | (R0) $\vdash \varphi, \varphi \rightarrow \psi \Rightarrow \vdash \psi$, |
| (A3) $K_a\varphi \rightarrow K_aK_a\varphi$, | (R1) $\vdash \varphi \Rightarrow \vdash K_a\varphi$, |
| (A4) $\neg K_a\varphi \rightarrow K_a\neg K_a\varphi$, | (R2) $\vdash \varphi \Rightarrow \vdash [\psi]\varphi$, |
| (A5) $[\varphi]p \leftrightarrow (\varphi \rightarrow p)$, | (R3) $\vdash \varphi \Rightarrow \vdash [G]\varphi$, |
| (A6) $[\varphi]\neg\psi \leftrightarrow (\varphi \rightarrow \neg[\varphi]\psi)$, | (R4) $\vdash \varphi \Rightarrow \vdash \langle[G]\rangle\varphi$, |
| (A7) $[\varphi](\psi \wedge \chi) \leftrightarrow ([\varphi]\psi \wedge [\varphi]\chi)$, | (R5) $(\forall\psi \in \mathcal{L}_{EL}^G \vdash \eta([\psi]\varphi)) \Rightarrow \vdash \eta([G]\varphi)$, |
| (A8) $[\varphi]K_a\psi \leftrightarrow (\varphi \rightarrow K_a[\varphi]\psi)$, | (R6) $(\forall\psi \in \mathcal{L}_{EL}^G \exists\chi \in \mathcal{L}_{EL}^{A \setminus G} \vdash \eta(\psi \rightarrow \langle\psi \wedge \chi\rangle\varphi)) \Rightarrow \vdash \eta(\langle[G]\rangle\varphi)$. |
| (A9) $[\varphi][\psi]\chi \leftrightarrow [\varphi \wedge [\varphi]\psi]\chi$, | |

So, **CoGAL** is the smallest subset of \mathcal{L}_{CoGAL} that contains all the axioms A0 – A11 and closed under rules of inference R0 – R6. Elements of **CoGAL** are called *theorems*. Note that R5 and R6 are infinitary rules: they require an infinite number of premises. Finding finite axiomatisations of any of **APAL**, **GAL**, or **CAL** is an open problem. Note also that **CoGAL** includes coalition logic [14], that is all the axioms of the latter are validities of **CoGAL** and a rule of inference preserves validity (see Appendix A).

Definition 2.6. (Soundness and completeness) An axiomatisation is *sound*, if for any formula φ of the language, it holds that $\varphi \in \mathbf{CoGAL}$ implies φ is valid. And vice versa for *completeness*.

Soundness of A0–A4, R0, and R1 is due to soundness of **S5**. Axioms A5–A9 and rule of inference R3 are sound, since **PAL** is sound [12]. Soundness of axiom A10 and rules of inference R3 and R5 was shown in [2]. We show soundness of R4, R6 in Proposition 2.7, and validity of A11 in Proposition 2.10.

Proposition 2.7. *R4 and R6 are sound, that is, they preserve validity.*

Proof. A proof is given in Appendix B (Proposition B.1). □

Validity of A11 was stated to be an open question in [11, 5]. Informally, the idea of our proof is as follows. Let us examine the axiom: $\langle[G]\rangle\varphi \rightarrow \langle G\rangle[A \setminus G]\varphi$. In the antecedent, all the agents make announcements simultaneously. In the consequent, the agents in $A \setminus G$ know the announcement ψ made by the agents in G . In the updated model (M^ψ, w) the agents in $A \setminus G$ may learn some *new* epistemic

formulae χ which they did not know before the announcement. We need to make sure that these new formulae cannot allow them to make φ false. However, since ψ is true in the initial model, and χ in the updated one, agents in $A \setminus G$ can always make an announcement that they know that after the announcement of ψ , χ holds. This announcement, made simultaneously with the announcement by G , ‘models’ the effect of announcing χ later. Returning to our example (Figure 1), whichever formulae ψ_1 and ψ_2 Ann and Bob announce, and whichever formula φ Cath learns afterwards, she can always announce $[\psi_1 \wedge \psi_2]K_c\varphi$ simultaneously with them in the initial situation. Informally, if after Bob’s announcement of $\neg p$, Cath learns that $\neg p$, she can announce: ‘If you say that $\neg p$ holds, then I will know it,’ or $[\neg p]K_c\neg p$. We use this idea to prove that if the agents in $A \setminus G$ can prevent φ after the announcement by G , then they could have prevented it before.

Due to restriction of announcements to formulas of epistemic logic, we cannot directly employ public announcement operators in agents’ ‘utterances.’ In order to avoid this, we use the standard translation of **PAL** into epistemic logic.

Definition 2.8. Translation function $t : \mathcal{L}_{PAL} \rightarrow \mathcal{L}_{EL}$ [12] is defined as follows:

$$\begin{aligned} t(p) &= p, & t([\varphi]\neg\psi) &= t(\varphi \rightarrow \neg[\varphi]\psi), \\ t(\neg\varphi) &= \neg t(\varphi), & t([\varphi](\psi \wedge \chi)) &= t([\varphi]\psi \wedge [\varphi]\chi), \\ t(\varphi \wedge \psi) &= t(\varphi) \wedge t(\psi), & t([\varphi]K_a\psi) &= t(\varphi \rightarrow K_a[\varphi]\psi), \\ t(K_a\varphi) &= K_at(\varphi), & t([\varphi][\psi]\chi) &= t([\varphi \wedge [\varphi]\psi]\chi). \\ t([\varphi]p) &= t(\varphi \rightarrow p), \end{aligned}$$

Every $\varphi \in \mathcal{L}_{PAL}$ is equivalent to $t(\varphi) \in \mathcal{L}_{EL}$.

Now we show that for every announcement of agents’ knowledge in some updated pointed model (M^ψ, w) there is an equivalent announcement in the original one (i.e. in (M, w)).

Lemma 2.9. Let $a, \dots, b \in A$. The following formula is valid for all $\psi, \chi_a, \dots, \chi_b \in \mathcal{L}_{EL}$:

$$[\psi \wedge K_at([\psi]\chi_a) \wedge \dots \wedge K_bt([\psi]\chi_b)]\varphi \leftrightarrow [\psi][K_a\chi_a \wedge \dots \wedge K_b\chi_b]\varphi$$

Proof. Suppose that for some pointed model (M, w) it holds that $(M, w) \models [\psi \wedge K_at([\psi]\chi_a) \wedge \dots \wedge K_bt([\psi]\chi_b)]\varphi$. By propositional reasoning, it is equivalent to $(M, w) \models [\psi \wedge (\psi \rightarrow K_at([\psi]\chi_a)) \wedge \dots \wedge (\psi \rightarrow K_bt([\psi]\chi_b))]\varphi$, and, by equivalence of a formula and its translation, the latter is equivalent to $(M, w) \models [\psi \wedge (\psi \rightarrow K_a[\psi]\chi_a) \wedge \dots \wedge (\psi \rightarrow K_b[\psi]\chi_b)]\varphi$. By A8, we have that $(M, w) \models [\psi \wedge [\psi]K_a\chi_a \wedge \dots \wedge [\psi]K_b\chi_b]\varphi$, and, by A7, $(M, w) \models [\psi \wedge [\psi](K_a\chi_a \wedge \dots \wedge K_b\chi_b)]\varphi$. Finally, by A9, the latter is equivalent to $(M, w) \models [\psi][K_a\chi_a \wedge \dots \wedge K_b\chi_b]\varphi$. \square

We use Lemma 2.9 to show validity of axiom A11.

Proposition 2.10. $\langle\!\langle G \rangle\!\rangle\varphi \rightarrow \langle G \rangle[A \setminus G]\varphi$ is valid.

Proof. Suppose to the contrary that for some pointed model (M, w) it holds that $(M, w) \models \langle\!\langle G \rangle\!\rangle\varphi$ and $(M, w) \not\models \langle G \rangle[A \setminus G]\varphi$. From $(M, w) \models \langle\!\langle G \rangle\!\rangle\varphi$, by the semantics,

$$\exists \psi \in \mathcal{L}_{EL}^G \forall \chi_a, \dots, \chi_b \in \mathcal{L}_{EL} : (M, w) \models \psi \wedge [\psi \wedge K_a\chi_a \wedge \dots \wedge K_b\chi_b]\varphi.$$

Let us call ψ_G the formula that G can announce to enforce φ . From $(M, w) \not\models \langle G \rangle[A \setminus G]\varphi$,

$$\forall \psi' \in \mathcal{L}_{EL}^G \exists \chi'_a, \dots, \chi'_b \in \mathcal{L}_{EL} : (M, w) \not\models \langle \psi' \rangle[K_a\chi'_a \wedge \dots \wedge K_b\chi'_b]\varphi.$$

In particular, for $\psi' = \psi_G$,

$$\exists \chi'_a, \dots, \chi'_b \in \mathcal{L}_{EL} : (M, w) \not\models \langle \psi_G \rangle [K_a \chi'_a \wedge \dots \wedge K_b \chi'_b] \phi.$$

Since ψ_G is true in (M, w) , this is equivalent to

$$\exists \chi'_a, \dots, \chi'_b \in \mathcal{L}_{EL} : (M, w) \not\models [\psi_G] [K_a \chi'_a \wedge \dots \wedge K_b \chi'_b] \phi.$$

By Lemma 2.9, the latter is equivalent to

$$\exists \chi'_a, \dots, \chi'_b \in \mathcal{L}_{EL} : (M, w) \not\models [\psi_G \wedge K_a t([\psi_G] \chi'_a) \wedge \dots \wedge K_b t([\psi_G] \chi'_b)] \phi.$$

Since $t([\psi_G] \chi'_a), \dots, t([\psi_G] \chi'_b)$ are in \mathcal{L}_{EL} , we have the contradiction with

$$\forall \chi_a, \dots, \chi_b \in \mathcal{L}_{EL} : (M, w) \models \psi_G \wedge [\psi_G \wedge K_a \chi_a \wedge \dots \wedge K_b \chi_b] \phi. \quad \square$$

Proposition 2.10 allows us to prove Lindenbaum Lemma (Proposition 2.19) for **CoGAL**. But before that, let us show some properties of the logic. The following validity shows that if some formula ϕ can be achieved by two coalition announcements, it can be achieved by a single joint coalition announcement as well. The validity was known only for the case of group announcements in **GAL** [2]. We show that this also holds for coalition announcements.

Proposition 2.11. $\langle [G] \rangle \langle [H] \rangle \phi \rightarrow \langle [G \cup H] \rangle \phi$ is valid.

Proof. The proof is presented in Appendix B (Proposition B.2). \square

Corollary 2.12. $\langle [G] \rangle \langle [G] \rangle \phi \rightarrow \langle [G] \rangle \phi$ is valid.

The other direction of Proposition 2.11 does not hold. Whether $\langle [G \cup H] \rangle \phi \rightarrow \langle [G] \rangle \langle [H] \rangle \phi$ is valid was posed as an open question in [5]. We settle this question by presenting a counterexample.

Proposition 2.13. $\langle [G \cup H] \rangle \phi \rightarrow \langle [G] \rangle \langle [H] \rangle \phi$ is not valid.

Proof. Let $G = \{a\}, H = \{b\}$, and $\phi := K_b(p \wedge q \wedge r) \wedge \neg K_a(p \wedge q \wedge r) \wedge \neg K_c(p \wedge q \wedge r)$. Informally, ϕ says that agent b knows that the given propositional variables are true, and agents a and c do not. Consider the model M in Figure 2 (reflexive arrows are omitted for convenience). By the semantics, $(M, \blacksquare) \models \langle \{a, b\} \rangle \phi$ iff $\exists (K_a \psi \wedge K_b \chi), \psi, \chi \in \mathcal{L}_{EL}, \forall K_c \tau, \tau \in \mathcal{L}_{EL} : (M, \blacksquare) \models K_a \psi \wedge K_b \chi \wedge [K_a \psi \wedge K_b \chi \wedge K_c \tau] \phi$. Let ψ be q , and χ be \top . Observe that $(M, \blacksquare) \models K_a q \wedge K_b \top$. Moreover, c does not know any formula that she can announce to avoid ϕ . An informal argument is as follows. Whatever c announces in this situation, she cannot avoid b learning $p \wedge q \wedge r$. In order to make a learn that $p \wedge q \wedge r$, c has to announce something of the form $\psi \rightarrow p$, since she does not know the value of p herself. Formula ψ can be neither r nor q , because c does not know their truth values. Also, it cannot be a statement of b 's knowledge, since in every q -world accessible by c , b 's knowledge is only a reflexive arrow. It cannot be a 's or c 's knowledge either, since in this case a would have known p herself, and c 's relation between q -states is universal.

In the consequent, we have $(M, \blacksquare) \models \langle [a] \rangle \langle [b] \rangle \phi$. By the semantics, $\exists \psi \in \mathcal{L}_{EL}^a \forall \chi \in \mathcal{L}_{EL}^{b \cup c} : (M, \blacksquare) \models \psi \wedge [\psi \wedge \chi] \langle [b] \rangle \phi$. Let us fix such a ψ , and let $\chi := K_b p \wedge K_c \top$. Then $(M, \blacksquare) \models \psi \wedge [\psi \wedge K_b p \wedge K_c \top] \langle [b] \rangle \phi$. Observe that no matter what a announces, $K_b p$ 'forces' a to learn that $p \wedge q \wedge r$, and whatever is announced in the updated model $(M^{\psi \wedge K_b p \wedge K_c \top}, \blacksquare)$, a 's knowledge of $p \wedge q \wedge r$ and, hence, falsity of ϕ remains. So, $(M, \blacksquare) \not\models \langle [a] \rangle \langle [b] \rangle \phi$. \square




Figure 2: A counterexample

2.3 Completeness

In order to prove completeness of **CoGAL**, we expand and modify the completeness proof for **APAL** [7, 9, 6]. Although the proof is partially based upon the classic canonical model approach, we have to ensure that construction of maximal consistent theories (Proposition 2.19) allows us to include infinite amount of formulae for cases of coalition announcements. This corresponds to ‘whatever other agents may say.’ But before we start, let us state two auxiliary lemmas.

Lemma 2.14. *Let $\varphi, \psi \in \mathcal{L}_{\text{CoGAL}}$. If $\varphi \rightarrow \psi$ is a theorem, then $\eta(\varphi) \rightarrow \eta(\psi)$ is a theorem as well.*

Proof. A simple induction on the complexity of η . □

Lemma 2.15. $[\chi]\langle\psi\rangle\varphi \rightarrow [\chi]\langle G\rangle\varphi$, where $\chi \in \mathcal{L}_{\text{CoGAL}}$, and $\psi \in \mathcal{L}_{\text{EL}}^G$ is a theorem.

Proof. Let $\psi \in \mathcal{L}_{\text{EL}}^G$. By A10, $[G]\neg\varphi \rightarrow [\psi]\neg\varphi$ is a theorem. By contraposition, we have that $\langle\psi\rangle\varphi \rightarrow \langle G\rangle\varphi$ is also a theorem. By R2 and distribution over implication, we infer $[\chi]\langle\psi\rangle\varphi \rightarrow [\chi]\langle G\rangle\varphi$. □

Now, the first part of the proof up to Proposition 2.19 is based on [7].

Definition 2.16. A set of formulae x is called a *theory* if and only if it contains **CoGAL**, and is closed under R0, R5, and R6. A theory x is consistent if and only if $\perp \notin x$, and is maximal if and only if for all $\varphi \in \mathcal{L}_{\text{CoGAL}}$ it holds that either $\varphi \in x$ or $\neg\varphi \in x$.

Proposition 2.17. *Let x be a theory, $\varphi, \psi \in \mathcal{L}_{\text{CoGAL}}$, and $a \in A$. The following are theories: $x + \varphi = \{\psi : \varphi \rightarrow \psi \in x\}$, $K_a x = \{\varphi : K_a \varphi \in x\}$, and $[\varphi]x = \{\psi : [\varphi]\psi \in x\}$.*

Proof. The proof is an extension of the one from [7] (see Appendix B, Proposition B.3). □

Proposition 2.18. *Let $\varphi \in \mathcal{L}_{\text{CoGAL}}$. Then, **CoGAL** + φ is consistent iff $\neg\varphi \notin \text{CoGAL}$.*

Proof. The proof is given in Appendix B (Proposition B.4). □

The following proposition is a variation of Lindenbaum Lemma. Validity of axiom A11 allows us to expand the corresponding proof for **APAL**, and to deal with having two different quantifiers at the same time.

Proposition 2.19. *Every consistent theory x can be extended to a maximal consistent theory y .*

Proof. Let ψ_0, ψ_1, \dots be an enumeration of formulae of the language, and let $y_0 = x$. Suppose that for some $n \geq 0$, y_n is a consistent theory, and $x \subseteq y_n$. If $y_n + \psi_n$ is consistent, then $y_{n+1} = y_n + \psi_n$. Otherwise, if ψ_n is not a conclusion of either R5 or R6, $y_{n+1} = y_n$. If ψ_n is a conclusion of R5, we enumerate all the subformulae of ψ_n which contain group announcement modalities $[G]$. Let $\eta_1([G]\varphi_1), \dots, \eta_k([G]\varphi_k)$ be all these subformulae. Then y_n^0, \dots, y_n^k is a sequence of consistent theories, where $y_n^0 = y_n$, and for some

$i < k$, y_n^i is a consistent theory containing y_n and $\neg\eta_i([G]\varphi_i)$. Since y_n^i is closed under $R5$, there exists $\chi \in \mathcal{L}_{EL}^G$ such that $\eta_i([\chi]\varphi_i) \notin y_n^i$. Hence, $y_n^{i+1} = y_n^i + \neg\eta_i([\chi]\varphi_i)$, and $y_{n+1} = y_n^k$.

Now, we consider the case when ψ_n is a conclusion of $R6$. We enumerate all the subformulae of ψ_n which contain coalition announcement modalities $[G]$. Let $\eta_1([G]\varphi_1), \dots, \eta_k([G]\varphi_k)$ be all these subformulae. Then y_n^0, \dots, y_n^k is a sequence of consistent theories, where $y_n^0 = y_n$, and for some $i < k$, y_n^i is a consistent theory containing y_n and $\neg\eta_i([G]\varphi_i)$. By $A11$, this means that $\neg\eta_i([G]\langle A \setminus G \rangle \varphi_i) \in y_n^i$. Since y_n^i is closed under $R5$, there exists $\chi \in \mathcal{L}_{EL}^G$ such that $\eta_i([\chi]\langle A \setminus G \rangle \varphi_i) \notin y_n^i$. Hence, $y_n^{i+1} = y_n^i + \neg\eta_i([\chi]\langle A \setminus G \rangle \varphi_i)$, and $y_{n+1} = y_n^k$. Note that since for all $\tau \in \mathcal{L}_{EL}^{A \setminus G}$ $\eta([\chi]\langle \tau \rangle \varphi) \rightarrow \eta([\chi]\langle A \setminus G \rangle \varphi)$ are theorems (by Lemmas 2.14 and 2.15), they and their contrapositions are already in y_n^i (since y_n^i is a theory). Thus, adding $\neg\eta_i([\chi]\langle A \setminus G \rangle \varphi_i)$ to y_n^i adds all the $\neg\eta_i([\chi]\langle \tau \rangle \varphi_i)$ for $\tau \in \mathcal{L}_{EL}^{A \setminus G}$ as well.

Finally, y is a maximal consistent theory, and $x \subseteq y$. \square

The rest of the proof is an expansion of the one from [9]. It employs induction on complexity of formulae to prove Truth Lemma (Proposition 2.25) and, ultimately, completeness (Proposition 2.26) of **CoGAL**.

Definition 2.20. The *size* of some formula $\varphi \in \mathcal{L}_{CoGAL}$ is defined as follows:

1. $Size(p) = 1$,
2. $Size(\neg\varphi) = Size(K_a\varphi) = Size([G]\varphi) = Size(\langle [G] \rangle \varphi) = Size(\varphi) + 1$,
3. $Size(\varphi \wedge \psi) = Size(\varphi) + Size(\psi) + 1$,
4. $Size([\psi]\varphi) = Size(\psi) + 3 \cdot Size(\varphi)$.

The \square -depth is defined as follows:

1. $d_{\square}(p) = 0$,
2. $d_{\square}(\neg\varphi) = d_{\square}(K_a\varphi) = d_{\square}(\langle [G] \rangle \varphi) = d_{\square}(\varphi)$,
3. $d_{\square}(\varphi \wedge \psi) = \max\{d_{\square}(\varphi), d_{\square}(\psi)\}$,
4. $d_{\square}([\psi]\varphi) = d_{\square}(\psi) + d_{\square}(\varphi)$,
5. $d_{\square}([G]\varphi) = d_{\square}(\varphi) + 1$.

The $\langle \diamond \rangle$ -depth is the same as \square , with the following exceptions.

1. $d_{\langle \diamond \rangle}([G]\varphi) = d_{\langle \diamond \rangle}(\varphi)$,
2. $d_{\langle \diamond \rangle}(\langle [G] \rangle \varphi) = d_{\langle \diamond \rangle}(\varphi) + 1$.

Definition 2.21. The binary relation $<_{\square, \langle \diamond \rangle}^{Size}$ between $\varphi, \psi \in \mathcal{L}_{CoGAL}$ is defined as follows: $\varphi <_{\square, \langle \diamond \rangle}^{Size} \psi$ iff $d_{\langle \diamond \rangle}(\varphi) < d_{\langle \diamond \rangle}(\psi)$, or, otherwise, $d_{\langle \diamond \rangle}(\varphi) = d_{\langle \diamond \rangle}(\psi)$, and either $d_{\square}(\varphi) < d_{\square}(\psi)$, or $d_{\square}(\varphi) = d_{\square}(\psi)$ and $Size(\varphi) < Size(\psi)$. The relation is a well-founded strict partial order between formulae.

Now, we ensure that the order of complexity is preserved. Case $[\bigwedge_{i \in G} K_i \psi_i] \varphi <_{\square, \langle \diamond \rangle}^{Size} [G] \varphi$ is obvious, since the public announcement on the left-hand side of the inequality is epistemic, and for any epistemic formula ψ , $d_{\square}(\psi) = d_{\langle \diamond \rangle}(\psi) = 0$. Case $[\chi][\bigwedge_{i \in G} K_i \psi_i] \varphi <_{\square, \langle \diamond \rangle}^{Size} [\chi][G] \varphi$ holds for the same reason. The cases for coalitions are identical: $\bigwedge_{i \in G} K_i \psi_i \rightarrow \langle \bigwedge_{i \in G} K_i \psi_i \wedge \bigwedge_{j \in A \setminus G} K_j \chi_j \rangle \varphi <_{\square, \langle \diamond \rangle}^{Size} \langle [G] \rangle \varphi$, and also $[\tau](\bigwedge_{i \in G} K_i \psi_i \rightarrow \langle \bigwedge_{i \in G} K_i \psi_i \wedge \bigwedge_{j \in A \setminus G} K_j \chi_j \rangle \varphi) <_{\square, \langle \diamond \rangle}^{Size} [\tau][G] \varphi$.

Definition 2.22. The *canonical model* is the model $M^C = (W^C, \sim^C, V^C)$, where

- W^C is the set of all maximal consistent theories,

- \sim^C is defined as $x \sim_a^C y$ iff $K_a x \subseteq y$,
- $x \in V^C$ iff $p \in x$.

Relation \sim^C is equivalence due to axioms A2, A3, and A4.

Definition 2.23. Let $\varphi \in \mathcal{L}_{CoGAL}$. Condition $P(\varphi)$: for all maximal consistent theories x , $\varphi \in x$ iff $x \in \llbracket \varphi \rrbracket_{M_C}$. Condition $H(\varphi)$: for all $\psi \in \mathcal{L}_{CoGAL}$, if $\psi <_{\llbracket \cdot, \{\cdot \} \rrbracket}^{Size} \varphi$, then $P(\psi)$.

Proposition 2.24. For all $\psi \in \mathcal{L}_{CoGAL}$, if $H(\varphi)$, then $P(\varphi)$.

Proof. Suppose $H(\varphi)$ holds, and let x be a maximal consistent theory. The proof is by induction on $<_{\llbracket \cdot, \{\cdot \} \rrbracket}^{Size}$ -complexity of formulae. Most of the cases were proved in [9]. We prove here only remaining instances involving group and coalition announcements.

Case $\varphi = [G]\psi$. Suppose that $[G]\psi \in x$. Since x is closed under R5, this is equivalent to $\forall \chi \in \mathcal{L}_{EL}^G$: $[\chi]\psi \in x$. By the fact that $[\chi]\psi <_{\llbracket \cdot, \{\cdot \} \rrbracket}^{Size} [G]\psi$, the latter holds if and only if $x \in \llbracket [\chi]\psi \rrbracket_{M_C}$ for all $\chi \in \mathcal{L}_{EL}^G$, which is equivalent to $x \in \llbracket [G]\psi \rrbracket_{M_C}$ by the semantics.

Case $\varphi = [\chi][G]\psi$. Suppose that $[\chi][G]\psi \in x$. Since $[\chi][G]\psi$ is a necessity form and x is closed under R5, this is equivalent to $\forall \tau \in \mathcal{L}_{EL}^G$: $[\chi][\tau]\psi \in x$. By the fact that $[\chi][\tau]\psi <_{\llbracket \cdot, \{\cdot \} \rrbracket}^{Size} [\chi][G]\psi$, the latter holds if and only if $x \in \llbracket [\chi][\tau]\psi \rrbracket_{M_C}$ for all $\tau \in \mathcal{L}_{EL}^G$, which is equivalent to $x \in \llbracket [\chi][G]\psi \rrbracket_{M_C}$ by the semantics.

Case $\varphi = \langle\langle G \rangle\rangle\psi$. Suppose that $\langle\langle G \rangle\rangle\psi \in x$. Since x is closed under R6, this is equivalent to $\forall \chi \in \mathcal{L}_{EL}^G$ $\exists \tau \in \mathcal{L}_{EL}^{A \setminus G}$: $\chi \rightarrow \langle \chi \wedge \tau \rangle \psi \in x$. By the fact that $\chi \rightarrow \langle \chi \wedge \tau \rangle \psi <_{\llbracket \cdot, \{\cdot \} \rrbracket}^{Size} \langle\langle G \rangle\rangle\psi$, the latter holds if and only if $\forall \chi \in \mathcal{L}_{EL}^G$ $\exists \tau \in \mathcal{L}_{EL}^{A \setminus G}$: $x \in \llbracket \chi \rightarrow \langle \chi \wedge \tau \rangle \psi \rrbracket_{M_C}$ which is equivalent to $x \in \llbracket \langle\langle G \rangle\rangle\psi \rrbracket_{M_C}$ by the semantics.

Case $\varphi = [\theta]\langle\langle G \rangle\rangle\psi$. Suppose that $[\theta]\langle\langle G \rangle\rangle\psi \in x$. Since $[\theta]\langle\langle G \rangle\rangle\psi \in x$ is a necessity form and x is closed under R6, this is equivalent to $\forall \chi \in \mathcal{L}_{EL}^G$ $\exists \tau \in \mathcal{L}_{EL}^{A \setminus G}$: $[\theta](\chi \rightarrow \langle \chi \wedge \tau \rangle \psi) \in x$. By the fact that $[\theta](\chi \rightarrow \langle \chi \wedge \tau \rangle \psi) <_{\llbracket \cdot, \{\cdot \} \rrbracket}^{Size} [\theta]\langle\langle G \rangle\rangle\psi$, the latter holds if and only if $\forall \chi \in \mathcal{L}_{EL}^G$ $\exists \tau \in \mathcal{L}_{EL}^{A \setminus G}$: $x \in \llbracket [\theta](\chi \rightarrow \langle \chi \wedge \tau \rangle \psi) \rrbracket_{M_C}$, which is equivalent to $x \in \llbracket [\theta]\langle\langle G \rangle\rangle\psi \rrbracket_{M_C}$ by the semantics. \square

Proposition 2.24 implies the following fact.

Proposition 2.25. Let $\varphi \in \mathcal{L}_{CoGAL}$, and x be a maximal consistent theory. Then $\varphi \in x$ iff $x \in \llbracket \varphi \rrbracket_{M_C}$.

Finally, we prove the completeness of **CoGAL**.

Proposition 2.26. For all $\varphi \in \mathcal{L}_{CoGAL}$, if φ is valid, then $\varphi \in \mathbf{CoGAL}$.

Proof. Towards a contradiction, suppose that φ is valid and $\varphi \notin \mathbf{CoGAL}$. Since **CoGAL** is a consistent theory, and by Propositions 2.17 and 2.18, we have that $\mathbf{CoGAL} + \neg\varphi$ is a consistent theory. Then, by Proposition 2.19, there exists a maximal consistent theory $x \supseteq \mathbf{CoGAL} + \neg\varphi$, such that $\neg\varphi \in x$. By Proposition 2.25, this means that $x \notin \llbracket \varphi \rrbracket_{M_C}$, which contradicts φ being a validity. \square

3 Conclusion

We presented **CoGAL** and provided a complete axiomatisation for it. The proof of completeness hinges on the validity of the axiom $\langle\langle G \rangle\rangle\varphi \rightarrow \langle G \rangle[A \setminus G]\varphi$. Validity of the other direction of the axiom, however, is still an open question. Answering it either way, positively, or negatively, will allow us to understand better mutual expressivity of **CAL** and **GAL**. The axiomatisation of **CoGAL** we presented is infinitary and employs necessity forms. Finding a finitary axiomatisation is yet another open problem. An interesting avenue of further research is adding common and distributed knowledge operators to **CoGAL** in the

vein of [1]. Additionally, since it is known that **GAL**, **CAL** [5], and hence **CoGAL**, are undecidable, a search for decidable fragments of these logics is another research question. We would also like to investigate applicability of logics with group and coalition announcements to epistemic planning [10]. Finally, a complete axiomatisation of **CAL** without group announcement operators has not been provided yet, and it is an intriguing direction of further research.

Acknowledgements

We would like to thank three anonymous reviewers for their insightful suggestions and detailed comments.

References

- [1] Thomas Ågotnes & Natasha Alechina (2012): *Epistemic coalition logic: completeness and complexity*. In: *International Conference on Autonomous Agents and Multiagent Systems, AAMAS 2012, Valencia, Spain, June 4-8, 2012 (3 Volumes)*, pp. 1099–1106. Available at <http://dl.acm.org/citation.cfm?id=2343854>.
- [2] Thomas Ågotnes, Philippe Balbiani, Hans van Ditmarsch & Pablo Seban (2010): *Group announcement logic*. *Journal of Applied Logic* 8(1), pp. 62–81, doi:10.1016/j.jal.2008.12.002.
- [3] Thomas Ågotnes & Hans van Ditmarsch (2008): *Coalitions and announcements*. In: *7th International Conference on Autonomous Agents and Multiagent Systems (AAMAS 2008), Estoril, Portugal, May 12-16, 2008, Volume 2*, pp. 673–680, doi:10.1145/1402298.1402318.
- [4] Thomas Ågotnes & Hans van Ditmarsch (2014): *Knowledge Games and Coalitional Abilities*. In: *Johan van Benthem on Logic and Information Dynamics*, Springer, pp. 451–485, doi:10.1007/978-3-319-06025-5_16.
- [5] Thomas Ågotnes, Hans van Ditmarsch & Timothy Stewart French (2016): *The Undecidability of Quantified Announcements*. *Studia Logica* 104(4), pp. 597–640, doi:10.1007/s11225-016-9657-0.
- [6] Philippe Balbiani (2015): *Putting right the wording and the proof of the Truth Lemma for APAL*. *Journal of Applied Non-Classical Logics* 25(1), pp. 2–19, doi:10.1080/11663081.2015.1011489.
- [7] Philippe Balbiani, Alexandru Baltag, Hans van Ditmarsch, Andreas Herzig, Tomohiro Hoshi & Tiago de Lima (2008): *‘Knowable’ as ‘known after an announcement’*. *Review of Symbolic Logic* 1(3), pp. 305–334, doi:10.1017/S1755020308080210.
- [8] Philippe Balbiani, Alexandru Baltag, Hans P. van Ditmarsch, Andreas Herzig, Tomohiro Hoshi & Tiago De Lima (2007): *What can we achieve by arbitrary announcements?: A dynamic take on Fitch’s knowability*. In: *Proceedings of the 11th Conference on Theoretical Aspects of Rationality and Knowledge (TARK-2007), Brussels, Belgium, June 25-27, 2007*, pp. 42–51, doi:10.1145/1324249.1324259.
- [9] Philippe Balbiani & Hans van Ditmarsch (2015): *A Simple Proof of the Completeness of APAL*. *Studies in Logic* 8(1), pp. 65–78. Available at <http://studiesinlogic.sysu.edu.cn:8080/ljxyj/EN/abstract/abstract211.shtml>.
- [10] Thomas Bolander & Mikkel Birkegaard Andersen (2011): *Epistemic planning for single and multi-agent systems*. *Journal of Applied Non-Classical Logics* 21(1), pp. 9–34, doi:10.3166/jancl.21.9-34.
- [11] Hans van Ditmarsch (2012): *Quantifying Notes*. In: *Logic, Language, Information and Computation - 19th International Workshop, WoLLIC 2012, Buenos Aires, Argentina, September 3-6, 2012. Proceedings*, pp. 89–109, doi:10.1007/978-3-642-32621-9_8.
- [12] Hans van Ditmarsch, Wiebe van der Hoek & Barteld Kooi (2008): *Dynamic Epistemic Logic*. *Synthese Library* 337, Springer, doi:10.1007/978-1-4020-5839-4.

- [13] Robert Goldblatt (1982): *Axiomatising the Logic of Computer Programming. Lecture Notes in Computer Science* 130, Springer, doi:10.1007/BFb0022481.
- [14] Marc Pauly (2002): *A Modal Logic for Coalitional Power in Games. Journal of Logic and Computation* 12(1), pp. 149–166, doi:10.1093/logcom/12.1.149.
- [15] Jan Plaza (2007): *Logics of public communications (reprint of 1989's paper). Synthese* 158(2), pp. 165–179, doi:10.1007/s11229-007-9168-7.
- [16] Bryan Renne, Joshua Sack & Audrey Yap (2009): *Dynamic Epistemic Temporal Logic. In: Logic, Rationality, and Interaction, Second International Workshop, LORI 2009, Chongqing, China, October 8-11, 2009. Proceedings*, pp. 263–277, doi:10.1007/978-3-642-04893-7_21.

A Coalition and Group Announcement Logic Subsumes Coalition Logic

Definition A.1. Axiomatisation of CL is as follows:

- | | |
|--|--|
| (C0) all instantiation of propositional tautologies, | (C4) $\langle\langle G \rangle\rangle(\varphi \wedge \psi) \rightarrow \langle\langle G \rangle\rangle\varphi$, |
| (C1) $\neg\langle\langle G \rangle\rangle\perp$, | (C5) $\langle\langle G \rangle\rangle\varphi \wedge \langle\langle H \rangle\rangle\psi \rightarrow \langle\langle G \cup H \rangle\rangle(\varphi \wedge \psi)$, |
| (C2) $\langle\langle G \rangle\rangle\top$, | if $G \cap H = \emptyset$, |
| (C3) $\neg\langle\langle \emptyset \rangle\rangle\neg\varphi \rightarrow \langle\langle A \rangle\rangle\varphi$, | (R0) $\vdash\varphi, \varphi \rightarrow \psi \Rightarrow \vdash\psi$, |
| | (R1) $\vdash\varphi \leftrightarrow \psi \Rightarrow \vdash\langle\langle G \rangle\rangle\varphi \leftrightarrow \langle\langle G \rangle\rangle\psi$. |

Proposition A.2. CoGAL contains CL.

Proof. C0 and R0 are already in CoGAL.

R1: $\vdash\varphi \leftrightarrow \psi \Rightarrow \vdash\langle\langle G \rangle\rangle\varphi \leftrightarrow \langle\langle G \rangle\rangle\psi$. Assume that $\vdash\varphi \leftrightarrow \psi$. This means that for any pointed model (M, w) the following holds: $(M, w) \models \varphi$ iff $(M, w) \models \psi$ (1). Now, suppose that for some pointed model (M, v) it holds that $(M, v) \models \langle\langle G \rangle\rangle\varphi$. By the semantics, $\exists\chi \in \mathcal{L}_{EL}^G \forall\tau \in \mathcal{L}_{EL}^{A \setminus G}: (M, v) \models \chi \wedge [\chi \wedge \tau]\varphi$, which is equivalent to the following: $(M, v) \models \chi$ and $((M, v) \models \chi \wedge \tau$ implies $(M^{\chi \wedge \tau}, v) \models \varphi$). By (1), we have that $\exists\chi \in \mathcal{L}_{EL}^G \forall\tau \in \mathcal{L}_{EL}^{A \setminus G}: (M, v) \models \chi$ and $((M, v) \models \chi \wedge \tau$ implies $(M^{\chi \wedge \tau}, v) \models \psi$), which is $(M, v) \models \langle\langle G \rangle\rangle\psi$ by the semantics. The same argumet holds in the other direction.

C1: $\neg\langle\langle G \rangle\rangle\perp$. From \top we derive $\langle\langle G \rangle\rangle\top$ by R4. Using the dual of box, we have $\neg\langle\langle G \rangle\rangle\neg\top$, or $\neg\langle\langle G \rangle\rangle\perp$.

C2: $\langle\langle G \rangle\rangle\top$. In any state, there exists a true announcement by G (each agent in G announces their knowledge of a tautology) and after any joint announcement, \top is true, hence, the axiom is valid.

C3: $\neg\langle\langle \emptyset \rangle\rangle\neg\varphi \rightarrow \langle\langle A \rangle\rangle\varphi$. By the semantics, $\neg\langle\langle \emptyset \rangle\rangle\neg\varphi$, which is $\langle\langle \emptyset \rangle\rangle\varphi$, means that there exists some $\psi \in \mathcal{L}_{EL}^A$, such that $\langle\psi\rangle\varphi$. This is precisely the meaning of $\langle\langle A \rangle\rangle\varphi$.

C4: $\langle\langle G \rangle\rangle(\varphi \wedge \psi) \rightarrow \langle\langle G \rangle\rangle\varphi$. Suppose that $\langle\langle G \rangle\rangle(\varphi \wedge \psi)$ holds. By the semantics, $\exists\chi \in \mathcal{L}_{EL}^G \forall\tau \in \mathcal{L}_{EL}^{A \setminus G}: \chi \wedge [\chi \wedge \tau](\varphi \wedge \psi)$. Then, by A7, we have $\exists\chi \in \mathcal{L}_{EL}^G \forall\tau \in \mathcal{L}_{EL}^{A \setminus G}: \chi \wedge [\chi \wedge \tau]\varphi \wedge [\chi \wedge \tau]\psi$. The latter implies $\exists\chi \in \mathcal{L}_{EL}^G \forall\tau \in \mathcal{L}_{EL}^{A \setminus G}: \chi \wedge [\chi \wedge \tau]\varphi$, which is $\langle\langle G \rangle\rangle\varphi$ by the semantics.

C5: $\langle\langle G \rangle\rangle\varphi_1 \wedge \langle\langle H \rangle\rangle\varphi_2 \rightarrow \langle\langle G \cup H \rangle\rangle(\varphi_1 \wedge \varphi_2)$, if $G \cap H = \emptyset$. Assume that $\langle\langle G \rangle\rangle\varphi_1 \wedge \langle\langle H \rangle\rangle\varphi_2$ holds. Let us consider the first conjunct. By the semantics, we have $\exists\psi_G \in \mathcal{L}_{EL}^G \forall\chi_{A \setminus G} \in \mathcal{L}_{EL}^{A \setminus G}: \psi_G \wedge [\psi_G \wedge \chi_{A \setminus G}]\varphi_1$ (1). The latter implies $\exists\psi_{G \cup H} \in \mathcal{L}_{EL}^{G \cup H} \forall\chi_{A \setminus G \cup H} \in \mathcal{L}_{EL}^{A \setminus G \cup H}: \psi_{G \cup H} \wedge [\psi_{G \cup H} \wedge \chi_{A \setminus G \cup H}]\varphi_1$. In order to show this let us assume to the contrary that $\forall\psi_{G \cup H} \in \mathcal{L}_{EL}^{G \cup H} \exists\chi_{A \setminus G \cup H} \in \mathcal{L}_{EL}^{A \setminus G \cup H}: \psi_{G \cup H} \rightarrow \langle\psi_{G \cup H} \wedge \chi_{A \setminus G \cup H}\rangle\neg\varphi_1$. Next, by (1), we fix some ψ_G . Then, we have $\forall\psi_H \in \mathcal{L}_{EL}^H \exists\chi_{A \setminus G \cup H} \in \mathcal{L}_{EL}^{A \setminus G \cup H}: \psi_G \rightarrow \langle\psi_G \wedge \psi_H \wedge \chi_{A \setminus G \cup H}\rangle\neg\varphi_1$, which means that there is a combination of ψ_H and $\chi_{A \setminus G \cup H}$ that makes φ_1 false. Since sets H and $A \setminus G \cup H$ comprise $A \setminus G$ (on condition that $G \cap H = \emptyset$), this means that there is some $\chi_{A \setminus G}$ which enforces $\neg\varphi_1$. Hence, the contradiction with (1). Similarly, we can prove that $\exists\psi_{G \cup H} \in \mathcal{L}_{EL}^{G \cup H}$

$\forall \chi_{A \setminus GUH} \in \mathcal{L}_{EL}^{A \setminus GUH}$: $\psi_{GUH} \wedge [\psi_{GUH} \wedge \chi_{A \setminus GUH}] \phi_2$. By distributivity of $[\]$ over \wedge , we get $\exists \psi_{GUH} \in \mathcal{L}_{EL}^{GUH}$
 $\forall \chi_{A \setminus GUH} \in \mathcal{L}_{EL}^{A \setminus GUH}$: $\psi_{GUH} \wedge [\psi_{GUH} \wedge \chi_{A \setminus GUH}] (\phi_1 \wedge \phi_2)$. Hence $\langle [G \cup H] \rangle (\phi_1 \wedge \phi_2)$. \square

B Proofs of Propositions

Proposition B.1. *R4 and R6 are sound, that is, they preserve validity.*

Proof. (R4) Assume that $\models \phi$. By R2, for an arbitrary ψ , $\models [\psi]\phi$. Since ψ is arbitrary, $\models \langle [G] \rangle \phi$; in other words, whatever agents announce, they cannot make a valid formula false.

(R6) Let (M, w) be an arbitrary pointed model. We proceed by induction on η .

Base case. $\forall \psi \in \mathcal{L}_{EL}^G \exists \chi \in \mathcal{L}_{EL}^{A \setminus G}$: $\psi \rightarrow \langle \psi \wedge \chi \rangle \phi$ is valid. Therefore, by the semantics, we infer validity of $\langle [G] \rangle \phi$.

Induction hypothesis. Assume the rule preserves validity for $n(\eta(\psi \rightarrow \langle \psi \wedge \chi \rangle \phi)) = k$. We show that it holds for $k + 1$.

Case $\forall \psi \in \mathcal{L}_{EL}^G \exists \chi \in \mathcal{L}_{EL}^{A \setminus G}$: $\tau \rightarrow \eta(\psi \rightarrow \langle \psi \wedge \chi \rangle \phi)$ is valid. This means that $(M, w) \models \tau \rightarrow \eta(\psi \rightarrow \langle \psi \wedge \chi \rangle \phi)$ iff $(M, w) \models \neg \tau$ or $(M, w) \models \eta(\psi \rightarrow \langle \psi \wedge \chi \rangle \phi)$, which is $(M, w) \models \neg \tau$ or $(M, w) \models \langle [G] \rangle \phi$ by Induction hypothesis. Hence, $(M, w) \models \tau \rightarrow \langle [G] \rangle \phi$.

Case $\forall \psi \in \mathcal{L}_{EL}^G \exists \chi \in \mathcal{L}_{EL}^{A \setminus G}$: $K_a \eta(\psi \rightarrow \langle \psi \wedge \chi \rangle \phi)$ is valid. This means that $(M, w) \models K_a \eta(\psi \rightarrow \langle \psi \wedge \chi \rangle \phi)$. By the semantics, for every $v \in W$: $(w, v) \in \sim_a$ implies $(M, v) \models \eta(\psi \rightarrow \langle \psi \wedge \chi \rangle \phi)$. By Induction hypothesis, for every $v \in W$: $(w, v) \in \sim_a$ implies $(M, v) \models \eta(\langle [G] \rangle \phi)$. And, by the semantics, $(M, w) \models K_a \eta(\langle [G] \rangle \phi)$.

Case $\forall \psi \in \mathcal{L}_{EL}^G \exists \chi \in \mathcal{L}_{EL}^{A \setminus G}$: $[\tau] \eta(\psi \rightarrow \langle \psi \wedge \chi \rangle \phi)$ is valid. This means that $(M, w) \models [\tau] \eta(\psi \rightarrow \langle \psi \wedge \chi \rangle \phi)$. By the semantics, $(M, w) \models \tau$ implies $(M^\tau, w) \models \eta(\psi \rightarrow \langle \psi \wedge \chi \rangle \phi)$. By Induction hypothesis, $(M, w) \models \tau$ implies $(M^\tau, w) \models \langle [G] \rangle \phi$. Finally, by the semantics, $(M, w) \models [\tau] \langle [G] \rangle \phi$. \square

Proposition B.2. $\langle [G] \rangle \langle [H] \rangle \phi \rightarrow \langle [G \cup H] \rangle \phi$ is valid.

Proof. Let $\psi := \bigwedge_{i \in G} K_i \psi_i$, $\psi' := \bigwedge_{j \in A \setminus G} K_j \psi'_j$, $\chi := \bigwedge_{k \in H} K_k \chi_k$, and $\chi' := \bigwedge_{l \in A \setminus H} K_l \chi'_l$. Suppose $(M, w) \models \langle [G] \rangle \langle [H] \rangle \phi$ for some M and $w \in W$. By the semantics,

$$\exists \psi \in \mathcal{L}_{EL}^G \forall \psi' \in \mathcal{L}_{EL}^{A \setminus G} \exists \chi \in \mathcal{L}_{EL}^H \forall \chi' \in \mathcal{L}_{EL}^{A \setminus H} : (M, w) \models \psi \wedge [\psi \wedge \psi'] (\chi \wedge [\chi \wedge \chi'] \phi).$$

By A7, we have: $(M, w) \models \psi \wedge [\psi \wedge \psi'] \chi \wedge [\psi \wedge \psi'] [\chi \wedge \chi'] \phi$. We are interested now in the third conjunct: $[\psi \wedge \psi'] [\chi \wedge \chi'] \phi$. By A9, we have that $(M, w) \models [\psi \wedge \psi' \wedge [\psi \wedge \psi'] (\chi \wedge \chi')] \phi$. Now, let us examine the following conjunction: $\exists \psi \in \mathcal{L}_{EL}^G \forall \psi' \in \mathcal{L}_{EL}^{A \setminus G}$: $\psi \wedge \psi'$, which is

$$\exists \bigwedge_{i \in G} K_i \psi_i \quad \forall \bigwedge_{j \in A \setminus G} K_j \psi'_j : \bigwedge_{i \in G} K_i \psi_i \wedge \bigwedge_{j \in A \setminus G} K_j \psi'_j$$

in the full form. We can present the set of agents $A \setminus G$ as a union of $G \cup H$ and $H \setminus G$ by expanding the right conjunct. So, we have $\exists \bigwedge_{i \in G} K_i \psi_i \quad \forall \bigwedge_{m \in H \setminus G} K_m \psi''_m \quad \forall \bigwedge_{j \in A \setminus GUH} K_j \psi'_j$:

$$\bigwedge_{i \in G} K_i \psi_i \wedge \bigwedge_{m \in H \setminus G} K_m \psi''_m \wedge \bigwedge_{j \in A \setminus GUH} K_j \psi'_j. \quad (1)$$

Since none of the universal quantifiers here is vacuous, there are particular ψ'' for which the conjunction holds. Formally, $\exists \bigwedge_{i \in G} K_i \psi_i \exists \bigwedge_{m \in H \setminus G} K_m \psi''_m \forall \bigwedge_{j \in A \setminus GUH} K_j \psi'_j$: (1). Therefore, combining G and $H \setminus G$, we have

$$\exists \bigwedge_{i \in GUH} K_i \psi_i \forall \bigwedge_{j \in A \setminus GUH} K_j \psi'_j : \bigwedge_{i \in GUH} K_i \psi_i \wedge \bigwedge_{j \in A \setminus GUH} K_j \psi'_j.$$

The same argument holds for the conjunction $\exists \chi \in \mathcal{L}_{EL}^H \forall \chi' \in \mathcal{L}_{EL}^G : \chi \wedge \chi'$. Let us redefine our auxiliary formulae: $\psi := \bigwedge_{i \in GUH} K_i \psi_i$, $\psi' := \bigwedge_{j \in A \setminus GUH} K_j \psi'_j$, $\chi := \bigwedge_{i \in GUH} K_i \chi_i$, and $\chi' := \bigwedge_{j \in A \setminus GUH} K_j \chi'_j$. Thus, we have that $\exists \psi \in \mathcal{L}_{EL}^{GUH} \forall \psi' \in \mathcal{L}_{EL}^{A \setminus GUH} \exists \chi \in \mathcal{L}_{EL}^{GUH} \forall \chi' \in \mathcal{L}_{EL}^{A \setminus GUH} : (M, w) \models [\psi \wedge \psi' \wedge (\psi \wedge \psi')(\chi \wedge \chi')]\varphi$. In the full form, the latter is

$$\begin{aligned} & \exists \bigwedge_{i \in GUH} K_i \psi_i \forall \bigwedge_{j \in A \setminus GUH} K_j \psi'_j \exists \bigwedge_{i \in GUH} K_i \chi_i \forall \bigwedge_{j \in A \setminus GUH} K_j \chi'_j : \\ & (M, w) \models [\bigwedge_{i \in GUH} K_i \psi_i \wedge \bigwedge_{j \in A \setminus GUH} K_j \psi'_j \wedge [\bigwedge_{i \in GUH} K_i \psi_i \wedge \bigwedge_{j \in A \setminus GUH} K_j \psi'_j] (\bigwedge_{i \in GUH} K_i \chi_i \wedge \bigwedge_{j \in A \setminus GUH} K_j \chi'_j)] \varphi. \end{aligned}$$

Using A7 and A8, we can ‘push’ announcements into the scope of knowledge operators:

$$\begin{aligned} (M, w) \models [\psi \wedge \psi' \wedge (\bigwedge_{i \in GUH} \bigwedge_{j \in A \setminus GUH} (K_i \psi_i \wedge K_j \psi'_j \rightarrow K_i [K_i \psi_i \wedge K_j \psi'_j] \chi_i) \wedge \\ \wedge (K_i \psi_i \wedge K_j \psi'_j \rightarrow K_j [K_i \psi_i \wedge K_j \psi'_j] \chi'_j))] \varphi. \end{aligned}$$

By propositional reasoning, the latter is equivalent to $(M, w) \models [\bigwedge_{i \in GUH} \bigwedge_{j \in A \setminus GUH} (K_i \psi_i \wedge K_j \psi'_j \wedge K_i [K_i \psi_i \wedge K_j \psi'_j] \chi_i \wedge K_j [K_i \psi_i \wedge K_j \psi'_j] \chi'_j)] \varphi$. Finally, we have

$$\begin{aligned} & \exists \bigwedge_{i \in GUH} K_i \psi_i \forall \bigwedge_{j \in A \setminus GUH} K_j \psi'_j \exists \bigwedge_{i \in GUH} K_i \chi_i \forall \bigwedge_{j \in A \setminus GUH} K_j \chi'_j : \\ & (M, w) \models [\bigwedge_{i \in GUH} \bigwedge_{j \in A \setminus GUH} (K_i (\psi_i \wedge [K_i \psi_i \wedge K_j \psi'_j] \chi_i) \wedge K_j (\psi'_j \wedge [K_i \psi_i \wedge K_j \psi'_j] \chi'_j))] \varphi. \end{aligned}$$

Conjuncts of the form $K_i(\psi_i \wedge [K_i \psi_i \wedge K_j \psi'_j] \chi_i)$ mean that agent i can announce ψ_i , i.e. what she knows now, or $[K_i \psi_i \wedge K_j \psi'_j] \chi_i$ (which is equivalent to $t([K_i \psi_i \wedge K_j \psi'_j] \chi_i)$), i.e. what she will know after announcements of other agents but not necessarily knows now, or both. Since all the variants comprise \mathcal{L}_{EL}^{GUH} , we rewrite the notation. Hence, $\exists \bigwedge_{i \in GUH} K_i \tau_i \forall \bigwedge_{j \in A \setminus GUH} K_j \tau'_j : (M, w) \models [\bigwedge_{i \in GUH} \bigwedge_{j \in A \setminus GUH} (K_i \tau_i \wedge K_j \tau'_j)] \varphi$, and at the same time $(M, w) \models \bigwedge_{i \in GUH} K_i \tau_i$ ($\bigwedge_{i \in G} K_i \psi_i$ is equivalent to $\bigwedge_{i \in GUH} K_i \psi_i$, where agents from H announce \top). And, by the semantics, this is $(M, w) \models \langle \langle GUH \rangle \rangle \varphi$. \square

Proposition B.3. *Let x be a theory, $\varphi, \psi \in \mathcal{L}_{CoGAL}$, and $a \in A$. The following are theories: $x + \varphi = \{ \psi : \varphi \rightarrow \psi \in x \}$, $K_a x = \{ \varphi : K_a \varphi \in x \}$, and $[\varphi]x = \{ \psi : [\varphi] \psi \in x \}$.*

Proof. We just expand the proof from [7] by showing that corresponding theories are closed under R5 and R6.

Suppose that $\eta([\psi]\chi) \in x + \varphi$ for all $\psi \in \mathcal{L}_{EL}^G$. It means that $\varphi \rightarrow \eta([\psi]\chi) \in x$ for all $\psi \in \mathcal{L}_{EL}^G$. Since $\varphi \rightarrow \eta([\psi]\chi)$ is a necessity form, and x is closed under R5, we infer that $\varphi \rightarrow \eta([G]\chi) \in x$, and, consequently, $\eta([G]\chi) \in x + \varphi$. So, $x + \varphi$ is closed under R5. Now, let $\forall \psi \in \mathcal{L}_{EL}^G \exists \tau \in \mathcal{L}_{EL}^{A \setminus G} : \eta(\psi \rightarrow \langle \psi \wedge \tau \rangle \chi) \in x + \varphi$. It means that $\forall \psi \in \mathcal{L}_{EL}^G \exists \tau \in \mathcal{L}_{EL}^{A \setminus G} : \varphi \rightarrow \eta(\psi \rightarrow \langle \psi \wedge \tau \rangle \chi) \in x$. Since

$\varphi \rightarrow \eta(\psi \rightarrow \langle \psi \wedge \tau \rangle \chi)$ is a necessity form, and x is closed under $R6$, we infer that $\varphi \rightarrow \eta(\langle [G] \rangle \chi) \in x$, and, consequently, $\eta(\langle [G] \rangle \chi) \in x + \varphi$. So, $x + \varphi$ is closed under $R6$.

Suppose that $\eta([\psi]\chi) \in K_a x$ for all $\psi \in \mathcal{L}_{EL}^G$. It means that $K_a \eta([\psi]\chi) \in x$ for all $\psi \in \mathcal{L}_{EL}^G$. Since $K_a \eta([\psi]\chi)$ is a necessity form, and x is closed under $R5$, we infer that $K_a \eta(\langle [G] \rangle \chi) \in x$, and, consequently, $\eta(\langle [G] \rangle \chi) \in K_a x$. So, $K_a x$ is closed under $R5$. Now, let $\forall \psi \in \mathcal{L}_{EL}^G \exists \tau \in \mathcal{L}_{EL}^{A \setminus G} : \eta(\psi \rightarrow \langle \psi \wedge \tau \rangle \chi) \in K_a x$. It means that $\forall \psi \in \mathcal{L}_{EL}^G \exists \tau \in \mathcal{L}_{EL}^{A \setminus G} : K_a \eta(\psi \rightarrow \langle \psi \wedge \tau \rangle \chi) \in x$. Since $K_a \eta(\psi \rightarrow \langle \psi \wedge \tau \rangle \chi)$ is a necessity form, and x is closed under $R6$, we infer that $K_a \eta(\langle [G] \rangle \chi) \in x$, and, consequently, $\eta(\langle [G] \rangle \chi) \in K_a x$. So, $K_a x$ is closed under $R6$.

Finally, suppose that $\eta([\psi]\chi) \in [\varphi]x$ for all $\psi \in \mathcal{L}_{EL}^G$. It means that $[\varphi]\eta([\psi]\chi) \in x$ for all $\psi \in \mathcal{L}_{EL}^G$. Since $[\varphi]\eta([\psi]\chi)$ is a necessity form, and x is closed under $R5$, we infer that $[\varphi]\eta(\langle [G] \rangle \chi) \in x$, and, consequently, $\eta(\langle [G] \rangle \chi) \in [\varphi]x$. So, $[\varphi]x$ is closed under $R5$. Now, let $\forall \psi \in \mathcal{L}_{EL}^G \exists \tau \in \mathcal{L}_{EL}^{A \setminus G} : \eta(\psi \rightarrow \langle \psi \wedge \tau \rangle \chi) \in [\varphi]x$. It means that $\forall \psi \in \mathcal{L}_{EL}^G \exists \tau \in \mathcal{L}_{EL}^{A \setminus G} : [\varphi]\eta(\psi \rightarrow \langle \psi \wedge \tau \rangle \chi) \in x$. Since $[\varphi]\eta(\psi \rightarrow \langle \psi \wedge \tau \rangle \chi)$ is a necessity form, and x is closed under $R6$, we infer that $[\varphi]\eta(\langle [G] \rangle \chi) \in x$, and, consequently, $\eta(\langle [G] \rangle \chi) \in [\varphi]x$. So, $[\varphi]x$ is closed under $R6$. \square

Proposition B.4. *Let $\varphi \in \mathcal{L}_{CoGAL}$. Then $\mathbf{CoGAL} + \varphi$ is consistent iff $\neg\varphi \notin \mathbf{CoGAL}$.*

Proof. From left to right. Suppose to the contrary that $\mathbf{CoGAL} + \varphi$ is consistent and $\neg\varphi \in \mathbf{CoGAL}$. Then, having both φ and $\neg\varphi$ means that $\perp \in \mathbf{CoGAL} + \varphi$, which contradicts to $\mathbf{CoGAL} + \varphi$ being consistent.

From right to left. Let us consider the contrapositive: if $\mathbf{CoGAL} + \varphi$ is inconsistent, then $\neg\varphi \in \mathbf{CoGAL}$. Since $\mathbf{CoGAL} + \varphi$ is inconsistent, $\perp \in \mathbf{CoGAL} + \varphi$, or, by Proposition B.3, $\varphi \rightarrow \perp \in \mathbf{CoGAL}$. By consistency of \mathbf{CoGAL} and propositional reasoning, we have that $\neg\varphi \in \mathbf{CoGAL}$. \square